Golden Eagle taken in West Florida.— An adult female Golden Eagle (Aquila chrysaëtos) was taken near DeFuniak Springs, Florida, on November 1, 1909. This is the third individual of this species taken in this vicinity during the past two years, all three specimens having been examined and identified by the writer. The first was taken on January 17, 1908, and the second on January 31, 1908.— G. CLYDE FISHER, DeFuniak Springs, Florida.

The Osprey a Breeder on the Catawba River, near Morganton, North Carolina.— Among the most noteworthy birds that were mounted and on exhibition in Leslie's drug store at Morganton, was a specimen of the Osprey (Pandion haliaëtus carolinensis). It was mounted with a Yellow-billed Cuckoo (Coccyzus americanus) in its talons! I was informed that the Osprey breeds regularly on the Catawba River, although I did not see any birds.

In his paper, 'An Ornithological Reconnaissance in Western North Carolina,' Mr. Brewster does not mention the Osprey. The country in the near vicinage of Morganton must be considered the westernmost breeding range for the species in the State.— Arthur T. Wayne, Mount Pleasant, S. C.

A New Name for Psephotus multicolor.— Psittacus multicolor (Temminek MS. 1819) "Brown" fide Kuhl, Conspectus Psittacorum, p. 55, No. 88 (1820); also Temminek, Trans. Linn. Soc. London, vol. 1, XIII, part 1, p. 119 (1821) (= Psephotus multicolor auctorum) is unfortunately preoccupied by Psittacus multicolor Gmelin, Syst. Nat., I, p. 328 (1788), applied to the Blue-bellied Lorikeet. The species commonly known as Psephotus multicolor may be called Psephotus varius.— Austin H. Clark, 1726 Eighteenth St., Washington, D. C.

Fork-tailed Flycatcher in Maine.— In late December, 1908, a correspondent wrote to me giving the description of a "peculiar bird" she had seen in the flesh while in the possession of an Indian guide and "taxidermist," and asking what it could possibly be. Her description made it absolutely certain that the bird must belong to the genus Muscivora, but which species was a matter requiring more data. After nearly a year's waiting I have at last had the specimen sent to me for identification and it is before me as this is written. It is a typical specimen of M. tyrannus, agreeing in measurements with those given by Ridgway for the female of the species. Though originally in good plumage it has been mounted by "main strength and foot power" and would require careful remounting to make it presentable.

The bird was shot near the mill of Mr. S. M. Holway in the town of